

***Aechmea distichantha* (Bromeliaceae) Epiphytes, Potential New Habitat for *Aedes Aegypti* and *Culex quinquefasciatus* (Diptera: Culicidae) Collected in the Province of Tucumán, Northwestern Argentina**

Author(s): Marina Stein María J. Dantur Juri , Griselda I. Oria and Patricia G. Ramirez

Source: Florida Entomologist, 96(3):1202-1206. 2013.

Published By: Florida Entomological Society

DOI: <http://dx.doi.org/10.1653/024.096.0368>

URL: <http://www.bioone.org/doi/full/10.1653/024.096.0368>

---

BioOne ([www.bioone.org](http://www.bioone.org)) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/page/terms\\_of\\_use](http://www.bioone.org/page/terms_of_use).

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

*AECHMEA DISTICHANTHA* (BROMELIACEAE) EPIPHYTES, POTENTIAL  
NEW HABITAT FOR *AEDES AEGYPTI* AND *CULEX QUINQUEFASCIATUS*  
(DIPTERA: CULICIDAE) COLLECTED IN THE PROVINCE OF TUCUMÁN,  
NORTHWESTERN ARGENTINA

MARINA STEIN<sup>1,\*</sup>, MARÍA J. DANTUR JURI<sup>2,3</sup>, GRISELDA I. ORIA<sup>1</sup> AND PATRICIA G. RAMIREZ<sup>1</sup>

<sup>1</sup>Instituto de Medicina Regional, Universidad Nacional del Nordeste, Avda. Las Heras 727, CP 3500, Resistencia, Chaco, Argentina

<sup>2</sup>Instituto Superior de Entomología "Dr. Abraham Willink", Facultad de Ciencias Naturales e Instituto Miguel Lillo, Universidad Nacional de Tucumán, Miguel Lillo 205, CP 4000 Tucumán, Argentina

<sup>3</sup>IAMRA, Universidad Nacional de Chilecito, 9 de Julio N° 22, CP F5360CKB, Chilecito, La Rioja, Argentina

Corresponding author; E-mail: marinastein@yahoo.com.ar

*Aedes* (*Stegomyia*) *aegypti* (Linnaeus) is a mosquito-associated with urban environments, and its larvae and pupae develop in a wide variety of artificial habitats (Forattini 1965). In the early 1960s, 17 countries and territories in the Americas, including Argentina, certified its eradication (Kerr et al. 1964). But a decade and a half later, they were re-colonized by *Ae. aegypti* due to the abandonment of the vector control programs. In 1987, the vector was again detected in Argentina, and during the early 1990s it colonized 16 provinces and 580 of the 854 existing municipalities (Coto & Masuh 2003). This is the only vector responsible for dengue epidemics that have occurred in our country from 1997 to date, recording more than 25,000 cases of DEN-1 in 2009 (Ministry of Public Health 2009).

Several studies conducted in Argentina report the presence of *Ae. aegypti* in a wide variety of artificial habitats in urban areas showing stable behavior of this species in relation to habitat preference, but also its ability to adapt to new conditions imposed by man (Campos 1993; Campos & Maciá 1996; Avilés et al. 1997; Schweigmann et al. 1997; Augier 1998; Almirón & Ludueña Almeida 1998; Stein & Oria 2002; Stein et al. 2002, 2007, 2011; Mangudo et al. 2010; Campos et al. 2011). In Brazil, *Ae. aegypti* has been collected from bromeliads in public and private gardens (Forattini et al. 2000; Mocellin et al. 2009), although as some authors suggested, these habitats do not contribute importantly to the production of its immatures (Mocellin et al. 2009; Santos et al. 2010). In relation to natural habitats in which *Ae. aegypti* has been observed in Argentina, Mangudo et al. (2010) found the species in treeholes in Aguaray city, Salta province, while Campos et al. (2011) observed this species in the axils of Araceae, at the National Park in Puerto Iguazú, province of Misiones. The phytotelmata have been identified as important habitats for vectors of human disease (O'Meara et al. 1995).

*Culex* (*Culex*) *quinquefasciatus* Say has also been found colonizing the same artificial locations as *Ae. aegypti* in urban environments, although it differs in behavior (Schweigmann et al. 1997; Almirón & Ludueña Almeida 1998; Oria et al. 1999, 2002; Stein & Oria 2002; Stein et al. 2002, 2007, 2011). This species is a competent vector of St. Louis Encephalitis virus in Argentina (Diaz 2009). Many other arboviruses have been isolated from *Cx. quinquefasciatus*, including Oropouche virus in Brazil. Vertical transmission of the Japanese encephalitis virus has also been demonstrated in this species (Lourenço-de-Oliveira & da Silva 1985; Johansen et al. 2001).

This is the first report of larvae of *Aedes aegypti* and *Culex quinquefasciatus* in epiphytic bromeliads in semi-urban and rural localities in the province of Tucumán, northwestern Argentina (Fig. 1). Both localities are in the subtropical mountainous rainforest also called Yungas that extends from the Bolivia border (S 22°) up to the north of Catamarca province (S 29°), traversing Salta, Jujuy and Tucumán provinces (Cabrera 1994; Brown & Grau 1995; Brown et al. 2001). In the Yungas it is possible to recognize 4 vegetation strata within forests (piedmont forest, mountain forest, and mountain wood and altitude grasses). The piedmont forest receives average annual precipitation of 820 mm (550-1400 mm) and the average annual temperature is 21.5 °C (average maximum = 27.6 °C, average minimum = 15.4 °C). The localities are situated in the piedmont forest, which is characterized by native vegetation patches with areas that are strongly modified by human activities (deforestation and agriculture). It is common to see increased plantings of sugar cane, soybean and citrus (Prado 1995; Dantur Juri et al. 2010a, 2010b). The forest is called "tipa and pacará" (*Tipuana tipu* (Benth.) Kuntze; Fabales: Fabaceae) and (*Enterolobium contortisiliquum* (Vell.) Morong; Fabales: Fabaceae), including canopy trees such as *Blepharocalyx salicifolius* (H.B.K.) (Myrtales: Myrtaceae),




Fig. 1. Map of the study area.

*Juglans australis* Griseb. (Fagales: Juglandaceae), *Terminalia triflora* (Griseb.) Lillo (Myrtales: Combretaceae), *Handroanthus impetiginosus* (Mart. ex DC.) Mattos (Lamiales: Bignoniaceae), *Jacaranda cuspidifolia* Mart. (Lamiales: Bignoniaceae), *Jacaranda mimosifolia* D. Don., *Tabebuia avellanedae* Lor. ex Griseb. (Lamiales: Bignoniaceae), *Tecoma stans* (L.) C. Juss. ex Kunth (Lamiales: Bignoniaceae), *Salix humboldtiana* Willd. (Malpighiales: Salicaceae), *Acacia visco* Griseb. (Fabales: Fabaceae), *Scutia buxifolia* Reissek (Rosales: Rhamnaceae), *Ceiba chodatii* Hassl. (Malvales: Malvaceae), *Celtis pubescens* (Kunth) Spreng. (Rosales: Cannabaceae), *Allophylus edulis* (St. Hil.) Radlkofler (Sapindales: Sapindaceae) and *Eugenia uniflora* L. (Myrtales: Myrtaceae), among others. Climber or aerial plant species belong to the families Bignoniaceae, Ulmaceae, and Amaranthaceae. There are also vascular epiphyte plants such as Bromeliaceae (Prado 1995; Dantur Juri et al. 2010a, 2010b).

Epiphytic bromeliads situated on various species of trees and on a roof of an abandoned construction were inspected in Monte Bello (S 27° 22' 43.89" W 65° 44' 30.02"; 535 m asl) and Iltico (S 27° 20' 13.56" W 65° 39' 25.07"; 411 m asl) localities (Fig. 1). They were found at heights between 0.50 m and up to 4 m above ground level. There were 2 collections, one in Nov 2012 and the other in Mar 2013 (Table 1, Fig. 2). We consider these sampling sites as natural habitats located in a residual primitive forest or wild ecosystem (Monte Bello) and the other as a primitive forest but modified by human activities (Iltico). The larvae and pupae were collected using a glass bottle (1,500

mL), with a rubber stopper with two holes through which were inserted 2 flexible plastic tubes. One tube was introduced into the bromeliad's water reservoir (phytotelm), and then by applying suction with the other tube, all the water contained in the phytotelm was collected together with the immature stages of the mosquitoes. The water containing the larvae and/or pupae was transported to the laboratory in jar labeled with names of collectors and the date and site of collection. The larvae and pupae were reared to obtain adults in the laboratory.

The culicids were sorted and identified following the Lane (1953) and Darsie (1985) determination keys. The bromeliads were also identified as *Aechmea distichantha* Lemaire (Poales: Bromeliaceae) a native species, in all cases. In Monte Bello we collected 5 larvae of *Ae. aegypti* and 1 larva and 1 pupa of *Cx. quinquefasciatus* from the bromeliads located on the roof of an abandoned building, only 10 m from the nearest house and the same distance from the primitive forest (Fig. 2a). At Iltico, we collected 6 larvae of *Ae. aegypti* from epiphytic bromeliads located 4 m high above ground on a tree at 2 m from the nearest houses (Fig. 2b). *Aedes aegypti* specimens were collected with larvae of *Culex* (*Culex*) *fernandezi* García & Cavalieri, *Culex* (*Microculex*) *imitator* Theobald and *Toxorhynchites* (*Lynchiella*) *guadalupensis* Dyar & Knab, while *Cx. quinquefasciatus* was collected with larvae of *Tx. guadalupensis* (Table 1).

*Aedes aegypti* was found in clear or turbid water, with or without organic matter, and with or without an unpleasant odor. *Culex quinquefas-*

TABLE 1. MOSQUITO SPECIES COLLECTED IN BROMELIADS FROM MONTE BELLO AND ILTICO LOCALITIES, PROVINCE OF TUCUMAN, BETWEEN NOV 2012 AND MAR 2013.

| Mosquito species                    | Localities  | Localities characteristics | Collection date | Number of specimens |
|-------------------------------------|-------------|----------------------------|-----------------|---------------------|
| <i>Aedes aegypti</i>                | Monte Bello | Primitive forest           | 13.IX.2012      | 3 larvae            |
|                                     | Monte Bello | Primitive forest           | 03.III.2013     | 2 larvae            |
|                                     | Iltico      | Modified forest            | 03.III.2013     | 6 larvae            |
| <i>Culex fernandezi</i>             | Monte Bello | Primitive forest           | 13.IX.2012      | 3 larvae            |
|                                     | Monte Bello | Primitive forest           | 03.III.2013     | 18 larvae           |
| <i>Culex imitator</i>               | Monte Bello | Primitive forest           | 13.IX.2012      | 28 larvae           |
|                                     | Monte Bello | Primitive forest           | 03.III.2013     | 15 larvae           |
|                                     | Iltico      | Modified forest            | 13.IX.2013      | 27 larvae           |
|                                     | Iltico      | Modified forest            | 03.III.2013     | 32 larvae           |
| <i>Culex quinquefasciatus</i>       | Monte Bello | Primitive forest           | 13.IX.2012      | 1 larva and 1 pupa  |
| <i>Toxorhynchites guadalupensis</i> | Monte Bello | Primitive forest           | 13.IX.2012      | 18 larvae           |
|                                     | Monte Bello | Primitive forest           | 03.III.2013     | 3 larvae            |
|                                     | Iltico      | Modified forest            | 13.IX.2013      | 10 larvae           |
|                                     | Iltico      | Modified forest            | 03.III.2013     | 5 larvae            |

*ciatus* was found in clear water without organic matter.

Despite the fact that the degree of infestation of nearby houses was not assessed either by the authors or by public health authorities, these observations should be taken into consideration by the latter, because the findings of *Ae. aegypti*

and *Cx. quinquefasciatus* in bromeliads located in wild and semi-urban environments could indicate a possible degree of invasion and adaptation to the primitive forest, or they may be a consequence of high infestations levels in nearby houses (Malta Verajao et al. 2005). This finding requires in-depth studies to reveal the real behavior of these



Fig. 2. Mosquito habitats provided by the *Aechmea distichantha*, a bromeliad epiphyte in the Yungas of north-western Argentina. (2a) Aspect of residual primitive forest in Monte Bello, meters from bromeliads where larvae of *Ae. aegypti* were found. (2b) Aspect of the abandoned building where epiphytic bromeliads were found from which the larvae of *Ae. aegypti* were collected. (2c) Epiphytic bromeliads from which larvae of *Ae. aegypti* and *Cx. quinquefasciatus* were collected at Iltico.



species in the study area, the true role of bromeliads as producers of these medically important mosquitos, and to determine and adopt appropriate control measures.

## SUMMARY

Larval habitats of *Aedes (Stegomyia) aegypti* (Linnaeus) and *Culex (Culex) quinquefasciatus* Say in the epiphyte *Aechmea distichantha* Lemaire (Poales: Bromeliaceae), were found and described both in semi-urban and rural localities of piedmont forest of the subtropical mountainous Yungas rainforest in the province of Tucumán, northwestern Argentina. This finding suggests that these anthropophilic disease vectors have achieved a degree of introduction and adaptation to the primitive forest, and that the bromeliad, which possesses phytotelmata, has an epidemiological role in providing natural water containers for the breeding of mosquito vectors.

Key Words: natural habitat, *Aedes aegypti*, *Culex quinquefasciatus*, phytotelmata, adaptation

## RESUMEN

*Aechmea distichantha* Lemaire (Poales: Bromeliaceae) epífita se describe como hábitat larval de *Aedes (Stegomyia) aegypti* (Linnaeus) y *Culex (Culex) quinquefasciatus* Say halladas tanto en localidades semi-urbanas como rurales de los bosques de piedemonte de la selva subtropical montañosa de las Yungas de la provincia de Tucumán, noroeste de Argentina. Este hallazgo sugiere que estos vectores antropofílicos han alcanzado cierto grado de introducción y adaptación al bosque primitivo, y que la bromelia (fitotelmata), tiene un papel epidemiológico en el suministro de contenedores naturales de agua para la cría de mosquitos vectores.

Palabras Clave: hábitat natural, *Aedes aegypti*, *Culex quinquefasciatus*, fitotelmata, adaptación

## REFERENCES CITED

- ALMIRÓN, W. R., AND BREWER, M. E. 1996. Classification of immature stage habitats of Culicidae (Diptera) collected in Córdoba, Argentina. *Mem Inst Oswaldo Cruz* 91: 1-9.
- ALMIRÓN, W. R., AND LUDUEÑA ALMEIDA, F. 1998. *Aedes aegypti* (Diptera: Culicidae) en Córdoba, Argentina. *Rev. Soc. Entomol. Argentina* 57: 27-28.
- AUGIER, L. M. 1998. Presencia de *Aedes aegypti* (Diptera: Culicidae) en Tucumán, Argentina. *Rev. Soc. Entomol. Argentina* 57: 66.
- AVILÉS, G., CECCHINI, R., HARRINGTON, M. E., CICHERO, J., AND ASIS, R. 1997. *Aedes aegypti* in Cordoba province, Argentina. *J. American Mosq. Control Assoc.* 13: 255-258.
- BROWN, D. A., AND GRAU, H. R. 1995. Investigación, conservación y desarrollo en las selvas subtropicales de montaña. Laboratorio de Investigaciones Ecológicas de las Yungas. Universidad Nacional de Tucumán. Tucumán, Argentina, 270 pp.
- BROWN, A. D., GRAU, H. R., MALIZIA, L., AND GRAU, A. 2001. Los Bosques Nublados de la Argentina, pp. 623-659 *In* M. Kappelle and A. D. Brown [ed.], *Bosques Nublados de Latinoamérica*, Editorial IN-Bio, Costa Rica.
- CABRERA, A. L. 1994. Regiones fitogeográficas argentinas. Tomo II. Enciclopedia Argentina de agricultura y jardinería. Buenos Aires. Editorial Acme S.A.C.I. 85 pp.
- CAMPOS, R. E. 1993. Presencia de *Aedes (Stegomyia) aegypti* L. (Diptera: Culicidae) en la localidad de Quilmes (Buenos Aires, Argentina). *Rev. Soc. Entomol. Argentina* 52: 36.
- CAMPOS, R. E., SPINELLI, G., AND MOGI, M. 2011. Culicidae and Ceratopogonidae (Diptera: Nematocera) inhabiting phytotelmata in Iguazú National Park, Misiones Province, subtropical Argentina. *Rev. Soc. Entomol. Argentina* 70: 111-118.
- COTO, H., AND MASUH, H. 2003. Control de *Aedes aegypti* (L.) en la ciudad de Clorinda (Formosa, Argentina): un modelo para áreas urbanas. *Actas VII Jornadas de Zoología del Uruguay*. pp. 140.
- DANTUR JURI, M. J., ZAIDENBERG, M., CLAPS, G. L., SANTANA, M., AND ALMIRÓN, W. R. 2010a. Abundance patterns of *Anopheles pseudopunctipennis* and *Anopheles argyritarsis* in northwestern Argentina. *Acta Trop.* 115: 234-241.
- DANTUR JURI, M. J., ALMIRÓN, W. R., AND CLAPS, G. L. 2010b. Population fluctuation of *Anopheles* (Diptera: Culicidae: Anophelinae) in natural and modified environments. *J. Vector Ecol.* 35: 28-34.
- DARSIE, R. F. JR. 1985. The mosquitoes of Argentina Parts I, Keys for identification of adult females and fourth stage larvae in English and Spanish (Diptera, Culicidae). *Mosq. Syst.* 17: 153-253.
- DIAZ, L. A. 2009. Patrones de actividad estacional del virus St. Louis encephalitis en Córdoba, Argentina. Tesis.UNC. 195 pp.
- FORATTINI, O. P. 1965. Entomología Médica. Vol. II. Culicini: *Culex*, *Aedes*, *Psorophora*. Ed. Univ. São Paulo, 506 pp.
- FORATTINI, O. P., AND ALVARENGA MONTEIRO MARQUES, G. R. 2000. Nota sobre o encontro de *Aedes aegypti* em bromélias *Rev. Saúde Pública* 34: 543-44.
- JOHANSEN, C. A., VAN DEN HURK, A. F., PYKE, A. T., ZBOROWSKI, P., PHILLIPS, D. A., MACKENZIE, J. S., AND RITCHIE, S. A. 2001. Entomological investigations of an outbreak of Japanese Encephalitis Virus in the Torres Strait, Australia, in 1998. *J. Med. Entomol.* 38: 581-88.
- KERR, J. A., CAMARGO, S. D., AND ABEDI, Z. H. 1964. Eradication of *Aedes aegypti* in Latin America. *J. American Mosq. Control Assoc.* 24: 276-82.
- LANE, J. 1953. Neotropical Culicidae, 1st/2nd ed., Ind Graf Siqueira SA, São Paulo, 1,112 pp.
- LOURENÇO DE OLIVEIRA, R., AND SILVA, T. F. 1985. Alguns aspectos da ecologia dos mosquitos (Diptera: Culicidae) de uma area de planície (Granjas Calabria), em Jacarepagua, Rio de Janeiro. III. Preferencia horaria das femeas para o hematofagismo. *Mem. Inst. Oswaldo Cruz* 80: 195-201.
- MALTA VAREJÃO, J. B., DOS SANTOS, C. B., REZENDE, H. R., BEVILACQUA, L. C., AND FALQUETO, A. 2005. Criadouros de *Aedes (Stegomyia) aegypti* (Linnaeus, 1762) em bromélias nativas na Cidade de Vitória, ES. *Rev. Soc. Brasil Med. Trop.* 38:238-240.

- MANGUDO, C., APARICIO, J. P., AND GLEISER, R. M. 2010. Huecos de árboles como hábitat larval de *Aedes aegypti* en el arbolado público de la localidad de Aguaray, provincia de Salta, VII Jornadas Regionales sobre mosquitos, Universidad Nacional de Misiones, Posadas, pp. 37.
- MINISTERIO DE SALUD, ARGENTINA 2009. In: <http://www.msal.gov.ar>
- MOCCELLIN, M. G., SIMÕES, T. C., FERNANDES SILVA DO NASCIMENTO, T., TEIXEIRA, M. L. F., LOUNIBOS, L. P., AND LOURENÇO DE OLIVEIRA, R. 2009. Bromeliad-inhabiting mosquitoes in an urban botanical garden of dengue endemic Rio de Janeiro. Are bromeliads productive habitats for the invasive vectors *Aedes aegypti* and *Aedes albopictus*? Mem. Inst. Oswaldo Cruz 104: 1171-1176.
- O'MEARA, G. F., EVANS, L. F., GETTMAN, A. D., AND PATTERSON, A. W. 1995. Exotic tank bromeliads harboring immature *Aedes albopictus* and *Aedes bahamensis* (Diptera: Culicidae) in Florida. J. Vector Ecol. 20: 216-24.
- ORIA, G. I., STEIN, M., AND GORODNER, J. O. 1999. Nuevos aportes en mosquitos (Diptera: Culicidae) de Resistencia, Chaco, Reunión de Comunicaciones Científicas y Tecnológicas de la UNNE, tomo VI, Ciencias Biológicas, Editorial Universitaria de la Universidad Nacional del Nordeste, Corrientes, pp. 31-34.
- ORIA, G. I., STEIN, M., AND GORODNER, J. O. 2002. Mosquitos, sus criaderos y factores socioculturales de la población en el nordeste argentino, pp. 167-172 In O. D. Solomón [ed.], Actualizaciones en Artropodología Sanitaria Argentina, Publicación Monográfica 2, Buenos Aires, Argentina.
- PRADO, D. E. 1995. Selva pedemontana: contexto regional y lista florística de un ecosistema en peligro, pp. 19-52 In A. D. Brown and H. R. Grau [eds.], Investigación, Conservación y Desarrollo en las selvas subtropicales de montaña, Laboratorio de Investigaciones Ecológicas de las Yungas, Universidad Nacional de Tucumán, Tucumán, Argentina.
- SANTOS, C. B., LEITE, G. R., AND FALQUETO, A. 2010. Do native bromeliads represent important breeding sites for *Aedes aegypti* (L.) (Diptera: Culicidae) in urbanized areas? Neotrop. Entomol. 40: 278-281.
- SCHWEIGMANN, N. J., VEZZANI, D., VERA, T., GÓMEZ, S., FERNÁNDEZ CAMPÓN, F., CEVASCO, C., FREIRE, G., KURUC, J., ACKERMANN, G., CARBAJO, A., BRUZZONE, O. A., BOFFI, R., AND ABRAMO ORREGO, L. 1997. Infestación domiciliar por formas inmaduras de *Aedes (Stegomyia) aegypti* L. En un foco del partido de San Martín, provincia de Buenos Aires, Argentina. Entomol Vectores. 4: 185-190.
- STEIN, M., AND ORIA, G. I. 2002. Identificación de criaderos de *Aedes aegypti* (Diptera: Culicidae) y cálculo de índices de infestación en la provincia del Chaco, pp. 161-166 In O. D. Solomón [ed.], Actualizaciones en Artropodología Sanitaria Argentina, Publicación Monográfica 2, Buenos Aires.
- STEIN, M., ORIA, G. I., AND ALMIRÓN, W. R. 2002. Principales criaderos para *Aedes aegypti* y culicidos asociados, Argentina. Rev. Saude Publica. 36: 627-630.
- STEIN, M., ALMIRÓN, W. R., LUDUEÑA ALMEIDA, F., ZAPATA, L., MARI, E., AND GORODNER, J. O. 2007. *Aedes aegypti* y culicidos asociados (Diptera: Culicidae) en la ciudad de Corrientes, Argentina. [www.siiisalud.com/dato/dat051/06d28002.htm](http://www.siiisalud.com/dato/dat051/06d28002.htm).
- STEIN, M., LUDUEÑA ALMEIDA, F., WILLENER, J. A., AND ALMIRÓN, W. R. 2011. Classification of immature mosquito species according to characteristics of the larval habitat in the subtropical province of Chaco, Argentina. Mem. Inst. Oswaldo Cruz 106: 400-407.